





Resection of a large portion of the liver is possible because the remaining liver grows to compensate for the part that was removed.

This determination can only be made by a surgeon experienced in liver surgery.


Liver resections: A, right hepatic lobectomy; B, left hepatic lobectomy; C, extended left hepatic lobectomy; D, extended right hepatic lobectomy; E, hepatic segmentectomy; F, non-anatomical wedge resection

Making the Determination for Liver Resection

Liver surgeons at the Johns Hopkins Medicine Liver Tumor Center are highly skilled in treating patients with advanced liver disease. In combination with other members of the multi-disciplinary team, your surgeon will assess if it is possible to remove the entire tumor or all of the disease while

leaving enough liver behind.

In some instances it may appear that the disease involves a large majority of your liver, and it may be determined that surgery is impossible. However, at our center, sometimes with the combination of chemotherapy to shrink the tumor and portal vein embolization to help the liver "grow" before the surgery, our surgeons are able to remove all the cancer in the liver.



Innovative Liver Surgery Techniques

chemotherapy is used to shrink the tumor in preparation for surgery.

Other innovative surgical techniques that have been widely adopted at Johns Hopkins Medicine include surgery to remove some of the masses combined with radio frequency ablation to destroy other masses with heat.

In some patients with extensive cancer in their liver, surgeons at Johns Hopkins Medicine have adopted a cutting-edge strategy of removal of all lesions in two operations - rather than one. Some patients would not have enough liver left after one operation. By using two operations, the liver is allowed to "grow back" after the first operation in order to allow for the second operation. This strategy may be used in patients who have significant tumors on both the right and left sides of the liver and who may not be considered surgical candidates at other institutions.

Minimally-Invasive Laparoscopic Liver Tumor Resection

For suitable patients, our surgeons also perform laparoscopic liver tumor removal which involves removing part of the liver with minimally invasive techniques, requiring a smaller incision and providing a quicker recovery.

- Home
- About Johns Hopkins Medicine
- Contact Johns Hopkins Medicine
- Centers & Departments
- Maps & Directions
- Find a Doctor
- **Patient Care**
- Education
- Research
- News
- Careers Site Map



Johns Hopkins Recent Recognitions





© The Johns Hopkins University, The Johns Hopkins Hospital, and Johns Hopkins Health System. All rights reserved.

Privacy Policy and Disclaimer